



A Comparative Study of Operative Fixation of Fracture of Medial Malleolus by Tension Band Wiring (TBW) & Cannulated Cancellus Screw (CC)

Aminul IM*

Orthopedics Department, Calcutta National Medical College, India

*Corresponding author: Aminul Islam Mamood, Assistant Professor, Orthopedics Department, Calcutta National Medical College, Kolkata, India, 700014, 7908543862, Email: aminulislamamamood@gmail.com

Received Date: February 09, 2024; Published Date: February 23, 2024

Abstract

Aim: To analyse the functional outcome of Ankle joint after medial malleolar fracture treated with ORIF with TBW and CC Screws.

Materials and Method: This is prospective study done in Calcutta national medical college and hospital, Kolkata from September 2019 to August 2020. In this study 20 cases of Medial malleolar ankle fracture were analysed. Muller Classification used. Road traffic accident, Twisting injury and fall from height are major mode of injury. TBW & 4.0 mm Cannulated Cancellous screw were used for fixation. Out of 20 cases 10 cases treated by TBW & 10 cases treated by 4mm CC screw.

Results: According to Baird and Jackson scoring system out of the 20 patients treated, Excellent and good functional results were achieved in 90% in TBW patients and in 80% in CCS patients with A better range-of-motion (especially for dorsiflexion of ankle) was noticed in TBW group. Radiological union took place in TBW patients (mean time 18 weeks) earlier than in CCS patients (mean time 20 weeks). On comparing the results in both groups, better results were seen in TBW group as compared to CCS group.

Conclusion: In this study we observed that the functional outcome of Ankle joint after medial malleolus fracture treated with open reduction and internal fixation with tension band wiring gives better results in range of motion and union more stable as compared to open reduction and internal fixation with 4.0 mm Cannulated Cancellous screws.

Keywords: Medial Malleolar Ankle Fracture; Muller Classification And Baird And Jackson Scoring System; Tension Band Wiring; Cannulated Cancellous Screw

Abbreviations: TBW: Tension Band Wiring; CCS: Cannulated Cancellus Screw; AO: Association of Osteosynthesis.

Introduction

Ankle joints are highly susceptible to injuries. This is because they are relatively mobile and bear much of the

stresses associated with weight bearing. The ankle joint supports more weight per unit area than any other joint in the body [1]. Ankle joint is the most congruous joint in the lower extremity bearing up to five times the body weight [2]. Sir Robert Jones said - Ankle is the most injured joint of the body but the least well treated [3]. Ankle injuries gain importance because body weight is transmitted through it and locomotion depends upon the stability of this joint. They

are usually mixed injuries, ligamentous and bony and each injury is an end result of the sequence of ligamentous and bony failure due to deforming forces.

The French have written extensively on ankle injuries beginning with Baron, Dupuytren and then Maisonneuve, but it was not until 1922 that a proper understanding of the mechanism and classification of the ankle fractures was published in a paper by Ashurst APC, et al. [4].

Fractures of the medial malleolus are one of the most common fractures encountered by an orthopaedic surgeon. Their incidence have increased over the last few decades & accounting for 9% of all fractures. It is difficult to restore the desired anatomical continuity and congruity of their articular surfaces after reduction. In the modern age, the treatment of osteoarticular fractures with plaster immobilisation for prolonged period is not entirely justified particularly when the fracture is sustained by relatively younger people.

Conservative treatment of these fractures with cast is becoming a past, early mobilization and early ambulation is the ideal treatment of fractures [5]. The superiority of ORIF over closed treatment have been thoroughly demonstrated in literature [6]. With evolution and advances in orthopaedics, the mode of treatment of osteoarticular fracture has changed from closed reduction and plaster immobilisation to open reduction and internal fixation. In treating medial malleolar fractures, restoration of anatomical alignment is highly essential. This is because, only a slight variation from normal is incompatible with good joint function. Ramsey and Hamilton have reported that only 1mm lateral shift of talus decreases contact area by 40% and displacement of 3 mm decreases contact area by 60 % [7].

Various implants made of stainless steel for fractures fixation, included screws, k-wires, tension band wires etc. Open reduction and internal fixation is an excellent method for restoration of normal anatomy of the joint. The study of Association of Osteosynthesis (AO) has evolved the methods of utilising rigid fixation and compression for internal fixation of fractures [5]. It works on principle that the implant absorbs tension and increases compression across fracture site and aims at full restoration of painless movement and strength.

Now a day's different implants and different modalities are available for the treatment of medial malleolar fractures, no method is perfect and each one has its advantages & disadvantages. So, aim of this study is to determine the outcome of medial malleolar fractures treated with open reduction and internal fixation with tension band wiring and 4 mm cannulated cancellous screws which will allow us to

practice more efficiently.

Materials & Methods

This study was done to evaluate and compare the outcomes of medial malleolar fractures operated with open reduction internal fixation with tension band wiring and 4mm cannulated cancellous screws in our hospital Calcutta National Medical College & Hospitals from September 2019 to august 2020. This is a prospective study of medial malleolar fractures in 20 patients treated at ours institution, the patients presenting themselves in casualty and OPD of department of orthopaedics and newly diagnosed as medial malleolar fractures. Follow up period was till 6 months. The institutional ethical committee clearance was taken before the start of the study.

When the patients were seen for the first time after injury, a through history was taken regarding the time of injury, mechanism of injury, any significant past or personal history. Patients were examined giving special importance to whether the fracture was open or closed, presence of gross swelling, fracture blisters and presence of other associated injuries.

Routine investigations were done as were necessary. The diagnosis was confirmed by antero-posterior, lateral and mortises radiographs. Stabilization was done with a below knee slab, and open reduction and fixation was done once the skin condition was good and swelling had subsided.

Inclusion Criteria

1. Patients of any sex.
2. Age group: 18 years above.
3. Patients who are fit for surgery. (No sever comorbidities)
4. Closed fractures
5. Transverse/Oblique vertical fracture.

Exclusion criteria

1. Open & commutated fractures
2. Patients unfit for surgery and/or anaesthesia. (Sever comorbidities)
3. Patients with history of previous medial malleolus fracture of either ankle.
4. Patients with vertical shear (adduction) fracture of medial malleolus.
5. Immature skeleton-Age below 18 years.

The form of treatment (tension band wiring or 4mm cannulated cancellous screws) was decided randomly.

Fractures type B, type C and type D according to Muller Classification for isolated fracture of medial malleolus were included in this study.

Muller classification for isolated fracture of medial malleolus

Type A: Fracture avulsions of the tip of the malleolus.

Type B: Fracture occurring between the tip and the level of the plafond

Type C: Fracture at the level of the plafond

Type D: Fracture extending vertically above this

Out of 20 fractures, 10 were operated with open reduction and internal fixation with tension band wiring and 10 were operated with open reduction and internal fixation with 4mm cannulated cancellous screws. The basic idea was to achieve near anatomical reduction. Open reduction and internal fixation was used in all 20 patients. It was done as a planned and elective procedure. Before patients were taken up for surgery, they were put on foot elevation and anti-inflammatory drugs for few days to reduce foot swelling. For all 20 cases anteromedial approach was used.

Post operatively limbs were immobilized in plaster splint, patients were put on antibiotics and analgesics and limbs were elevated. Dressings were done regularly and sutures were removed on average on 14th day (decided according to the wound inspection). The below knee splint was continued till 6 weeks post operatively in both groups and was then removed. Patients were assessed for fracture union radiologically. Once the fracture showed signs of union, partial weight bearing was advised, which was approximately at 6-8 weeks and it was then gradually increased to full weight bearing. Physiotherapy exercises for ankle movements were started on removal of slab.

Follow up

Weight bearing is restricted for 6 weeks. At 6 weeks the plaster was removed. Clinical examination was done regarding movement of ankle. At 6 weeks x-ray of the ankle was taken in AP, lateral and mortise views and looked for signs of fracture union and then were advised partial weight bearing once the fracture showed signs of union. Patients were advised to keep the limb at elevation at night times and perform active movements of ankle joints. It was then gradually increased to full weight bearing. Regular follow up was done at 6 weeks, 3 months and thereafter at monthly intervals till 6 months after operation till the fracture united. Follow up X-rays were taken to assess fracture union, the condition of implant (Figures 1 & 2). Union was considered when fracture line could not be seen on radiographs and normal bony trabecular were seen crossing across the

fracture site. When patient could walk full weight bearing and did not have any tenderness.



Figure 1(a): Pre-Operative X-ray.

Radiological outcome of ankle joint after tension band wiring fixation



Figure 1(b): Post-operative X-ray.

Radiological outcome of Ankle joint after CC-screw Fixation



Figure 2(a): Pre-Operative X-ray.




Figure 2(b): Post-operative X-ray.

Assessment of Results

Baird and Jackson scoring system was used to evaluate the patients at the end of 6 months. Score and fracture union in both the groups were compared using unpaired-t|| test. Finally correlation between results of the score and clinical data were evaluated using chi square test.

Baird and Jackson scoring system

1) Pain Score

No Pain	15
Mild pain with strenuous activity	12
Mild pain with activities of daily living	8
Pain with weight bearing	4
Pain at rest	0

2) Stability of ankle

No clinical instability	15
Instability with sports activities	5
Instability with activities of daily living ability to walk	0

3) Able to walk

Able to walk desired distances without limp or pain	15
Able to walk desired distances with mild limp or pain	12
Moderately restricted in ability to walk	8
Able to walk short distances only	4
Unable to walk	0

4) Able to run

Able to run desired distances without pain	10
Able to run desired distances with slight pain	8
Moderate restriction in ability to run with mild pain	6
Able to run short distances only	3
Unable to run	0

5) Ability to work

Able to perform usual occupation without restrictions	10
Able to perform usual occupation with restrictions in some strenuous activities	8
Able to perform usual occupation with substantial restriction	6
Partially disabled; selected jobs only	3
Unable to work	0

6) Motion of the ankle

Within 10 of uninjured ankle	10
Within 15 of uninjured ankle	7
Within 20 of uninjured ankle	4
<50 of uninjured ankle, or dorsiflexion <5	0

7) Radiographic result

Anatomical with intact mortise (normal medial clear space, Normal 2mm superior joint space, no talar tilt)	25
Same as above with mild reactive changes at the joint margins	15
Measurable narrowing of the superior joint space, superior joint space 2mm, or talar tilt >2mm	10
Moderate narrowing of the superior joint space, with superior space between 2 and 1mm.	5
Severe narrowing of the superior joint space, with superior joint space <1mm widening of the medial clear space, severe reactive changes (Sclerotic subchondral bone and osteophyte formation)	0

A scores according to the Baird and Jackson Scoring system

Excellent	96-100
Good	91-95
Fair	81-90
Poor	0-80
Maximum possible score	100

Observation and Results

A total of 20 cases were enrolled in the study, 10 cases were operated with open reduction and internal fixation with 4mm cannulated cancellous screws and 10 cases were operated with open reduction and internal fixation with tension band wiring. The mean age in CCS group was 45 years whereas in TBW group was 48 years (Table 1).

Majority of the patients (60 %) included in the study were in the age group 40-59 years. When Chi square test was applied for the distribution of patients in three age groups for both the operative procedures modalities it was found to be statistically non-significant. Out of total 20 patients considered in study 15 (75%) were male and 5 (25%) were female. Almost similar ratio was seen in either of the groups. Out of total 20 patients treated 8 (40%) had medial malleolus fracture on right side and 12 (60%) had medial malleolus fracture on left side. The comparison of side affected in individual group was statistically non-significant.

In CC-Screw group, 4 (40%) patients had Type B fracture, 4 (40%) patients had Type C fracture and 2 (20%) patients had Type D fracture. In TBW group, 2 (20%) patient had Type B fracture, 5 (50%) patients had Type C fracture and 3 (30%) patients had Type D fracture (Table 2).

At the end of 6 weeks, the mean range of dorsiflexion in CCS groups was 13 as compare to 15.66 in TBW group, was not statistically significant and the mean range of plantar flexion in CCS group was 18.66 degrees as compared to 20.33 degrees in TBW group, which was not statistically significant. At the end of 3 months, the mean range of dorsiflexion in CCS group was 17 degrees as compared to 19 degrees in TBW group, which was not statistically significant and the mean range of plantar flexion in CCS group was 23 degrees as compared to 25 degrees range of plantar flexion in TBW group. The improvement in the range was statistically significant (p value- 0.0274) in TBW group.

At the end of 6 months, the mean range of dorsiflexion in CCS group was 24.5 degrees as compared to 28 degrees in TBW group and showed no statistically significant difference (Figure 1). The mean range of plantar flexion in CCS group was 37 degrees as compared to 39.5 degrees range of plantar flexion in TBW group (Figure 2). The improvement in the range was statistically significant in TBW group with a p value of 0.0439.

Study Group	Mean Age
CCS	45
TBW	48

Table 1: Mean Age.

Study Group	Type B	Type C	Type D
CCS	4 (40%)	4 (40%)	2(20%)
TBW	2 (20%)	5(50%)	3(30%)

Table 2: No & Type of Fractures.




Figure 1: After 6 month mean range of dorsiflexion.




Figure 2: After 6 month mean range of planter flexion.

Out of total 20 patients treated in this study, 3 cases in CCS group and 2 cases in TBW group developed postoperative infection which was similar in both groups and was statistically insignificant.

At the end of 6 months, Average time required for radiological union in CCS group was 20 weeks and in TBW group was 18 weeks (Figure 3). Union occurred earlier in TBW group. When unpaired t test was applied to these values, it was found to be statistically significant with a p value 0.0303.




Figure 3: Time taken for radiological union.

The mean Baird and Jackson score at the end of 6 months for CCS group was 93.40 and for TBW group was 94.70 the mean score in TBW group was found to be better as compared to CCS group. . On applying unpaired t test, statistically significant difference was found (p value- 0.018). Out of the 20 patients treated 45% had excellent results, 40% had good results and 15% had fair results (Figure 4). On comparing the results in both groups by applying chi square test statistically significant difference was found with p value of 0.0490, thus results of TBW groups had better results than CCS groups.




Figure 4: Baird-Jackson score at the end of 6 months.

Out of the 10 patients in CC-Screw group Type B fracture Treated had 30% good results and 10% fair result, Type C fracture had 20% good results and 20% had fair results and Type D fracture had 20% Excellent results with the mean Baird and Jackson score at the end of 6 months for Type B- 91, for Type C- 89 and for Type D- 97 (Figure 5).




Figure 5: Results on the Basis of fracture Type CCS group.

Out of the 10 patients in TBW group Type B fracture Treated had 20% good results, Type C fracture had 30% good results and 20% Fair results Type D fracture had 30% Excellent results with the mean Baird and Jackson score at the end of 6 months for Type B- 93, for Type C-90 and for Type D- 98 (Figure 6).




Figure 6: Results on the Basis of fracture Type TBW group.

Discussion

Malleolar injuries are the most common significant lower extremity fracture. They primarily involve medial malleolus. They are produced indirectly by shearing and tensile forces applied through the talus [8,9]. The treatment of medial malleolar fractures with accurate open reduction and stable internal fixation using AO method and principles was found to give a high percentage of excellent and good results. In this study the outcomes of medial malleolar fractures treated with open reduction and internal fixation with tension band wiring and 4 mm cannulated cancellous screws were evaluated and compared. The results of present study are compared to with those of previous similar studies.

The mean age of this study was 46.50 years. This finding was similar to observation of Roberts RS [10] whose study the mean age was 40 years, Beris AE, et al. [11] in whose study the mean age was 43.8 years. There was male predominance in this series which is also observed in other studies. In this study out of 20 patients, 15(75%) were male and 05(25%) were females. Shams N, et al. [12] in his study of 30 patients had 22 male and 8 female patients. Maruthi CV, et al. [13] et al in their study of 40 cases had 28 males and 12 females.

In this study we observed that 40% are excellent and 40% are good results in CCS group (treated with 4mm cannulated cancellous screws) and 50% are excellent and 40% are good results in TBW group (treated with tension-band wiring) according to Baird and Jackson scoring system . This agrees with the results of Sang-Hanko M, et al. [14] who achieved excellent and good results in about 78% of cases treated with screws and 89% of cases treated with tension-band wiring. Al-Lamy WA, et al. [15] also achieved excellent and good results in 80% in group treated with screws and 90% in group treated with tension-band wiring, according to the modified ankle scoring system of Olerud C, et al. [16] (Table 3).

Study	CCS (Excellent to Good)	TBW (Excellent to Good)
Sang- Hanco & Young-Junpark [14]	78%	89%
Al-Lamy et al. [15]	80%	90%
Present Study	80%	90%

Table 3: Results in various studies.

In this study the mean time for radiological bone union in CCS group was 20 weeks and in TBW group was 18 weeks. So union occurred earlier in tension band wiring group in this study. This is similar to Nurul SK, et al. [17] study that reported a mean time of 11.8 weeks for screws and 9.4 weeks for tension-band wiring; and Al-Lamy WA, et al. [15] who in their study had union time of 12 weeks in screw group and 9 weeks in tension band wiring group, both of whom reported earlier fracture union in tension band wiring group (Table 4).

Study	CCS (Weeks)	TBW (Weeks)
SK.Nurul et al. [17]	11.8	9.4
Al-Obaidy and Al-Lamy [15]	12	9
Present Study	20	18

Table 4: Radiological Union in various studies.

Ostrum RF, et al. [18] recently demonstrated the biomechanics advantages of the tension-band over other fixation techniques for medial malleolus. When resisting pronation forces and applying compression force tension-band were four times stronger than screws. This might explain the faster union rate we were achieved in TBW group (mean of 18 weeks) as compared with CCS group (mean of 20 weeks).

We did not have any cases of non-union which were similar to the study results of Nurul SK, et al. [17] who achieved 100% union rate in both groups. Al-Lamy WA, et al. [15] had experienced two cases of delayed union (5%) out of 40 cases of the study but did not report any cases of non-union. Mack D, Szabo RM study reported loss of reduction with the use of tension band technique as a result of K-wires become loosened and migrate proximally [19]. Some reported back out of cannulated cancellous screws leading to implant loosening. On the other hand many authors did not agree with the frequency of this complication and reported that with the proper surgical techniques, wire migration was not a problem [20, 21].

In this study we did not see any wire migration, pull out of screw or loss of reduction and implant failure. Al-Lamy

WA, et al. [15] in their study of 40 cases did not report any case of fixation failure or kirschner wires migration or loss of reduction. Better range of motion was noticed in TBW group as compared with CCS group. This could be attributed to more rigid fixation with the use of tension-band. The use of tension band wire for fixation on the tension surface of a fractured bone converts the distracting tensile force in to a compressive force [22].

In this study out of 20 cases, 3 cases in CCS & 2 cases TBW group, postoperative complication we encountered was superficial infection, which was managed by regular dressings and administration of antibiotics. Shams N, et al. [12] in his study of 30 patients, reported superficial infection in 2 patients. Choudhari P, et al. [23] in their study on 68 cases, superficial infection was found in only one patient.

Conclusion

In this study we observed that the functional outcome of Ankle joint after medial malleolus fracture treated with open reduction and internal fixation with tension band wiring gives better results in range of motion and union as compared to open reduction and internal fixation with 4.0 mm cannulated cancellous screws. Thus, we conclude that open reduction and internal fixation with tension band wiring give better functional and radiological outcomes as compared to cannulated cancellous screws.

Funding

No funding sources.

Conflict of Interest

None declared.

Ethical approval

The study was approved by the institutional ethics committee.

References

1. Roy I, Davidovitch EAK (2009) Ankle Fractures. In: Bucholz WR, Charles, Heckman JD, Paultornetta, (Eds.), Rockwood and Greens Fractures in Adults. 7th (Edn.), Philadelphia: Lippincott Williams and Wilkins Publishers pp: 1975-2017.
2. Simon WH, Friedenburg S, Richardson S (1973) Joint congruence. A Correlation of Joint Congruence and Thickness of Articular Cartilage in Dogs. J Bone Joint Surg Am 55(8): 1614-1620.
3. Marvin SL (1994) Complication of Fractures and Dislocation of the Ankle. In: EPPS, Charles H (Eds.),

- Complications in Orthopaedic Surgery 3rd (Edn.), JB Lippincott Company, Philadelphia 1: 595-648.
4. Ashurst APC, Bromer RS (1992) Classification and Mechanism of Fracture of Leg Bones Involving the Ankle: Based on a Study of 300 Cases from the Episcopal Hospital. Arch Surg 4(1): 51-129.
 5. Muller ME, Allgower M, Scheinder R, Willenegger H (1995) Manual of Internal Fixation: Techniques Recommended by AO Group. 3rd (Edn.), Springer Verlag, New York 147: 148-149.
 6. Weber MJ (1993) Ankle Fractures and Dislocations. In: Chapman MW, Philadelphia MM (Eds.), Operative orthopaedics. 2nd (Edn.), J B Lippincott Company 3: 731-748.
 7. Ramsey L, Hamilton W (1987) Changes in Tibiotalar Area of Contact Caused by Lateral Talar Shift. J Bone Joint Surg Am 58(3): 356-357.
 8. Daly PJ, Fitzgerald RH, Melton LJ, Ilstrup DM (1987) Epidemiology of Ankle Fractures in Rochester, Minnesota. Acta Orthop Scand 58(5): 539-544.
 9. Lauge-Hansen N (1950) Fractures of the Ankle. II. Combined Experimental-Surgical and Experimental-Roentgenologic Investigations. Arch Surg 60(5): 957-985.
 10. Roberts RS (1983) Surgical Treatment of Displaced Ankle Fractures. Clin Orthop Relat 172: 164-170.
 11. Beris AE, Kabbani KT, Xenakis TA, Mitsionis G, Soucacos PK, et al. Surgical Treatment of Malleolar Fractures. A Review Of 144 Patients. Clin Orthop Related Research 341: 90-98.
 12. Shams N, Ahmed I, Hegde A (2014) A Study on Surgical Treatment of Ankle Fractures- A Clinical Study of 21 Cases. IJBAR 5(4): 190-192.
 13. Maruthi CV, Venugopal N, Nanjundappa HC, Siddalingaswamy MK (2014) Bimalleolar Fracture of Ankle Joint Managed by Tension Band Wiring Technique: A Prospective Study. Sch J App Med Sci 2(1D): 428-432.
 14. Sang-Hank M, Young-Jun PD (2002) Comparison between Screw Fixation and Tension-Band Wiring for Medial Malleolus Fractures. Korean Society of Foot Surg 6(1): 41-44.
 15. Al-Lamy WA, Al-Obaidy MH (2008) Comparative Study of Internal Fixation for Displaced Closed Fracture of Medial Malleolus Using Malleolar Screw versus Tension-band Wiring. Karbala J Med 2(4-5): 308-318.
 16. Olerud C, Molander H (1984) A Scoring Scale for Symptom Evaluation after Ankle Fracture. Arch Orthop Trauma Surg 103: 190-194.
 17. Nurul SK, Parviz Shahidi (1998) Comparative Study of Malleolar Fractures by Tension-Band and Malleolar Screw. BOS Journal 12(1): 13-19.
 18. Ostrum RF, Listsky AS (1997) Tension-Band Fixation of Medial Malleolar Fracture. J Orthop Trauma 6 (4): 464-468.
 19. Mack D, Szabo RM (1985) Complications of Tension-Band Wiring of Olecranon Fractures. J Bone Joint Surg 67(9): 1396-1401.
 20. Kinik H, Us AK, Mergen E (2002) Self-Locking Tension Band Technique. Arch Orthop Trauma Surg 119: 432-434.
 21. Joy G, Patzakis MJ, Harvey JP (1974) Precise Evaluation of the Reduction of Severe Ankle Fractures. J Bone Joint Surg 56(5): 979-993.
 22. AO (2017) Principles of fracture fixation 1: 249-254.
 23. Choudhari P, Agrawal R (2015) Comparative Study of One Versus Two Screws in Treatment of Isolated Fracture of Medial Malleolus. International Journal of Current Medical and Applied Sciences 7(3): 151-154.